



Annual Review of Marine Science

Marine Heatwaves

Eric C.J. Oliver,¹ Jessica A. Benthuyssen,²
Sofia Darmaraki,¹ Markus G. Donat,³
Alistair J. Hobday,⁴ Neil J. Holbrook,^{5,6}
Robert W. Schlegel,⁷ and Alex Sen Gupta^{8,9}

¹Department of Oceanography, Dalhousie University, Halifax, Nova Scotia B3H 4R2, Canada;
email: eric.oliver@dal.ca

²Australian Institute of Marine Science, Crawley, Western Australia 6009, Australia

³Barcelona Supercomputing Center, 08034 Barcelona, Spain

⁴CSIRO Oceans and Atmosphere, Hobart, Tasmania 7000, Australia

⁵Institute for Marine and Antarctic Studies, University of Tasmania, Hobart, Tasmania 7001, Australia

⁶Australian Research Council Centre of Excellence for Climate Extremes, University of Tasmania, Hobart, Tasmania 7001, Australia

⁷Department of Physical Oceanography, Woods Hole Oceanographic Institution, Woods Hole, Massachusetts 02543, USA

⁸Climate Change Research Centre, The University of New South Wales, Sydney, New South Wales 2052, Australia

⁹Australian Research Council Centre of Excellence for Climate Extremes, The University of New South Wales, Sydney, New South Wales 2052, Australia

Annu. Rev. Mar. Sci. 2021. 13:20.1–20.30

The *Annual Review of Marine Science* is online at
marine.annualreviews.org

<https://doi.org/10.1146/annurev-marine-032720-095144>

Copyright © 2021 by Annual Reviews.
All rights reserved

Keywords

climate extremes, sea surface temperature, climate change, mixed-layer dynamics, ocean circulation, air–sea heat flux, climate variability

Abstract

Ocean temperature variability is a fundamental component of the Earth's climate system, and extremes in this variability affect the health of marine ecosystems around the world. The study of marine heatwaves has emerged as a rapidly growing field of research, given notable extreme warm-water events that have occurred against a background trend of global ocean warming. This review summarizes the latest physical and statistical understanding of marine heatwaves based on how they are identified, defined, characterized, and monitored through remotely sensed and in situ data sets. We describe the physical mechanisms that cause marine heatwaves, along with their global distribution, variability, and trends. Finally, we discuss current issues in this developing research area, including considerations related to the



choice of climatological baseline periods in defining extremes and how to communicate findings in the context of societal needs.

1. INTRODUCTION

Marine heatwaves (MHWs)—a term first used by Pearce et al. (2011)—are anomalous warm sea-water events that can substantially affect marine ecosystems. An MHW is qualitatively defined as a discrete period of prolonged anomalously warm water at a particular location, and quantitative definitions are based on ocean temperatures exceeding a fixed (Frölicher et al. 2018), seasonally varying (Hobday et al. 2016), or cumulative (Eakin et al. 2010) threshold. MHWs are caused by a combination of local oceanic and atmospheric processes, including air–sea heat flux and horizontal temperature advection, and may be modulated by large-scale climate variability, including remote sources via teleconnections (Holbrook et al. 2019). MHWs have been identified primarily using ocean surface temperatures, although they may also extend below the surface (Schaeffer & Roughan 2017). The global count of MHW days per year has risen over the historical record due to increases in MHW duration and frequency (Oliver et al. 2018a). This trend is projected to increase further under climate change (Frölicher et al. 2018, Oliver et al. 2019) as a consequence of long-term ocean warming.

The study of MHWs as discrete events is motivated largely by their ecological and socioeconomic impacts. The study of sea surface temperature (SST) variability has a long history in physical oceanography and climate science (Philander 1983, Deser et al. 2010), and understanding the effects of warm ocean temperature extremes on marine ecosystems is vitally important (Garrahou et al. 2009, Mills et al. 2013, Wernberg et al. 2013, Smale et al. 2019, Benthuyssen et al. 2020). Specifically, studies of MHWs are set apart from studies of SST variability more broadly in that they examine the aspects of SST variability that affect marine ecosystems, rather than just the physical and climatic properties of ocean temperature variability.

Here, we critically review our current understanding of MHWs from a physical and climatological perspective. We review the definitions of MHWs, data sets useful for monitoring them, the physical processes causing MHWs, and their statistical properties. We examine the global distribution of MHWs, including characteristics of their variability and trends, and the role of anthropogenic change. The field of MHWs has grown as events continue to occur that have broad impacts across marine ecosystems and socioeconomic systems, raising issues of how best to communicate about MHWs and the most appropriate way to define MHWs in a warming ocean.

2. A BRIEF HISTORY OF RECENT HIGH-IMPACT EVENTS

While anomalously warm seawater events are not a new phenomenon, they have occurred with increasing frequency and duration over the past century (Oliver et al. 2018a) and have had significant impacts on marine ecosystems. The global-scale coral bleaching events of 1998, 2010, and most recently 2014–2017 occurred with El Niño conditions and globally elevated ocean temperatures (Eakin et al. 2019). The increased occurrence of extreme warm-water events across the globe has focused efforts toward examining events within a common framework. Over the past two decades, analyses of prominent MHW events (**Figure 1**) have revealed that they are often caused by compounding influences of atmospheric forcing and/or oceanic processes. Furthermore, the phase of climate modes of variability can act to increase or decrease the likelihood of MHW occurrence regionally (Holbrook et al. 2019). Here, we highlight notable events that have been attributed to local changes in air–sea heat fluxes, ocean heat transport, or remote forcing via teleconnections.

Marine heatwave (MHW): a discrete period of prolonged anomalously warm water at a particular location

Threshold: a temperature value, which may be fixed or vary over time and/or space, above which temperatures are considered anomalously warm in defining MHWs

Air–sea heat flux: turbulent (sensible and latent) and radiative (longwave and shortwave) exchanges of heat between the ocean and atmosphere

Temperature advection: a local temperature change due to the action of ocean currents in the presence of a spatial gradient in temperature, which may act in the horizontal (e.g., geostrophic boundary currents, wind-driven Ekman flows) or the vertical (e.g., upwelling/downwelling)

MHW duration: the time from the start date to the end date of an MHW




Figure 1

Key historical MHWs. (a) SST anomalies (above 1°C) on the day of peak MHW intensity. MHW intensity was defined based on the time series of SST averaged over the regions indicated by the black boxes. Light gray indicates areas of sea ice influence. (b) MHW properties for key historical events. The MHW intensity (y axis), MHW duration (x axis), and category (color; see Hobday et al. 2018a) were determined from the spatially averaged time series, as in panel a. The MHW area (circle size) is the total contiguous area reaching at least category 2 (strong). All events shown in panel b are referenced in Section 2. Abbreviations: AL, Gulf of Alaska and Bering Sea; Beng., Benguela; ECS, East China Sea; Med., Mediterranean; NA, northern Australia; MHW, marine heatwave; NEP, northeast Pacific; NWA, northwest Atlantic; SST, sea surface temperature; Tas., Tasman Sea; WA, Western Australia; WSA, western South Atlantic. Panel a inspired by a schematic from Frölicher & Laufkötter (2018).

Record-breaking MHWs have been documented where the atmospheric state played a central role in their development and maintenance. Examples of such events include those in the Mediterranean Sea during the summers of 2003 (Olita et al. 2007) and 2006 (Mavrakakis et al. 2018), off the northeast coast of the United States in 2012 (Chen et al. 2014, 2015), from the southeast tropical Indian Ocean to the Coral Sea in 2015–2016 (Benthuisen et al. 2018), in the East China Sea in

Climate modes of variability: recurrent spatial patterns of specific climate variables, often quasi-periodic or broad-banded and occurring on subseasonal, interannual, decadal, multidecadal, centennial, millennial, or longer timescales

MHW intensity: the temperature anomaly (based on the maximum or mean value) during an MHW

Anomaly: a deviation of temperature from the climatological mean

2016 (Tan & Cai 2018), along the southwest Atlantic shelf in 2017 (Manta et al. 2018), off coastal Peru in 2017 (Echevin et al. 2018), and in the Tasman Sea in 2017–2018 (Perkins-Kirkpatrick et al. 2019, Salinger et al. 2019). In these cases, the anomalously warm ocean temperatures were related to abnormally high air–sea heat fluxes into the ocean.

Other MHWs have been found to be caused primarily by anomalous ocean heat transport, such as the MHWs in the Angola Benguela Upwelling System in 2001 (Rouault et al. 2007), off Western Australia in early 2011 (Feng et al. 2013, Benthuyssen et al. 2014), in the Tasman Sea in 2015–2016 (Oliver et al. 2017), and in the Middle Atlantic Bight (Gawarkiewicz et al. 2019) and off Japan (Sugimoto et al. 2020) in 2017. In addition, large-scale MHWs have developed through a combination of multiple interacting mechanisms. Most notably, the 2014–2016 MHW affected the Pacific and was linked to tropical–extratropical teleconnections (Bond et al. 2015, Di Lorenzo & Mantua 2016, Hu et al. 2017, Tseng et al. 2017). Other events have been associated with substantial retreat in Arctic sea ice (e.g., the Gulf of Alaska and the Bering Sea in 2016; Walsh et al. 2018, Oliver et al. 2018b). There is an increasing interest in such events that may span across systems, including the ocean, atmosphere, land, and cryosphere (e.g., Ruthrof et al. 2018).

While these recent MHWs had varying intensities, durations, and spatial extents (**Figure 1b**), they have all been linked with disruptive changes in marine ecosystems (Wernberg et al. 2016, Hughes et al. 2017, Smale et al. 2019) and commercial fisheries (Mills et al. 2013, Caputi et al. 2016, Jacox et al. 2019). It should be noted that MHWs are distinct from otherwise anomalously warm years. Smale et al. (2019) pointed out that, while most recent research has been on the ecological effects of mean climate variables, discrete events have emerged as critical factors driving rapid shifts in ecosystems. These issues highlight the need for consistent MHW detection methods, frameworks for understanding their physical drivers, and accurate dynamical and statistical models for predicting future events.

3. DEFINING AND MONITORING MARINE HEATWAVES

3.1. Defining a Marine Heatwave

Extreme events are climate anomalies that are rare and have magnitudes that deviate significantly from typical conditions in a region. The statistics of extremes, those that describe the tails of a probability distribution, have traditionally been examined using extreme value theory (Gumbel 1958, Leadbetter et al. 1983, Coles 2001). Extreme value theory examines the distribution of extreme values in a time series by assuming the data are distributed according to certain probability distributions, such as the generalized extreme value (for block-maxima extremes, e.g., the annual maxima in **Figure 2**) or the generalized Pareto distribution (for peak-overthreshold extremes). The fitted distribution can be used to estimate the return periods or return levels of extreme events. Extreme value theory has not been widely applied to study temperature extremes in the ocean (e.g., Oliver et al. 2014a,b).

More recently, methods for detecting ocean temperature extremes have used a threshold above which any contiguous days of temperature are considered to be a single event. In this analysis, the important choice is in defining the threshold, either fixed in time or allowed to vary seasonally or on longer timescales. Fixed thresholds can be defined by an absolute temperature (**Figure 2a**), informed by marine species' upper thermal limits (the temperatures above which biological functions become impaired). For example, cumulative temperatures above an absolute threshold have been used for coral bleaching (e.g., degree heating days, weeks, and months; Liu et al. 2014). Temporally fixed thresholds can be defined using a relative measure of the temperature variance (e.g., quantiles). While these thresholds are fixed in time, similar to the absolute thresholds, this



Figure 2 (Figure appears on preceding page)

Different methods of detecting extreme temperatures for the 2011 Western Australia MHW (Pearce & Feng 2013) (*left column*) and 2012 northwest Atlantic MHW (Mills et al. 2013) (*right column*). (a) Fixed thresholds—defined either by an absolute temperature (here set at 25°C for Western Australia and 17°C for the northwest Atlantic; *dashed green lines*) or a fixed percentile threshold (here the 90th percentile; *solid green lines*)—and annual maximum temperature values (*white circles*) work well for detecting events during the warm season (as in Western Australia in 2011) but are poor indicators of anomalous conditions during other seasons (as in the northwest Atlantic in 2012). (b) Seasonally varying percentile thresholds, such as the Hobday et al. (2016) definition using the 90th percentile, allow detection of MHWs as anomalously warm temperatures during warm and cool seasons (including both the 2011 Western Australia MHW and the 2012 northwest Atlantic MHW). (c) Increasing the threshold—for example, by using the 99th percentile instead of the 90th—isolates only the most extreme events. (d) The categorization scheme, proposed by Hobday et al. (2018a), permits the use of a lower percentile threshold while identifying category levels of increasing intensity. The climatology was calculated using NOAA OI SST V2.0 data with a baseline period of 30 years (1983–2012). Abbreviations: MHW, marine heatwave; NOAA OI SST V2.0, National Oceanic and Atmospheric Administration Optimum Interpolation Sea Surface Temperature version 2.0.

Climatology: the statistical properties of a time series defined over an extended period (e.g., 30 years), including the mean, seasonal cycle, variance, and quantiles

Baseline period: the time period used to calculate a climatology, traditionally taken to be a 30-year period, following World Meteorological Organization recommendations

measure allows the definition of extreme to come from the time series data distribution itself (**Figure 2a**). This approach may be less applicable to a single species, but the results represent a more general quantification of what is extreme in the local context. Fixed thresholds typically only identify warm-season MHWs (e.g., coral bleaching thresholds; Fordyce et al. 2019). However, extremely warm temperatures in other seasons may have important consequences for the life cycle and survival of marine species (e.g., the survival of invasive warm-water species in winter; Ling et al. 2009). Temperatures below a lower threshold are known as marine cold spells (see the sidebar titled Marine Cold Spells).

A more versatile method to identify MHWs uses a seasonally varying climatological threshold (**Figure 2b**). The climatology is calculated over a chosen baseline period, preferably a 30-year temperature time series (WMO 2018). The method defined by Hobday et al. (2016) uses a local daily upper-percentile climatology as the threshold above which MHWs are detected, consistent with atmospheric heatwave definitions (Perkins & Alexander 2013). A 90th-percentile threshold results in a relatively large number of MHW events, and some weak, short-lived events may not result in ecological consequences. One solution is to use a higher threshold, such as the 99th percentile (Frölicher et al. 2018, Collins et al. 2019) (**Figure 2c**). This method identifies more intense MHWs but requires a longer observational record to estimate the threshold in a robust way, which can be problematic given the limitations of many observational records (see Section 3.2).

MARINE COLD SPELLS

Prolonged ocean temperatures in the lower tail of the temperature distribution are marine cold spells (MCSs) and may be defined analogously to MHWs by their exceedance of temperatures below a percentile threshold for a prolonged period (Schlegel et al. 2017). MCSs can be driven by intense upwelling or atmospheric forcing (e.g., cold-air outbreaks or storm-induced mixing). Severe MCSs can cause fish kills and invertebrate mortality and can pose risks to coral reefs (Gunter 1951, Roberts et al. 1982, Holt & Hold 1983, Firth et al. 2011, Paz-García et al. 2011). Higher-latitude systems have some resilience (Tuckett & Wernberg 2018), and MCSs may have historically inhibited the tropicalization of temperate ecosystems (Lerriorato & Nakamura 2019). MCSs have also been implicated as a controlling factor in ecosystem shifts between salt marshes and mangroves (Cavanaugh et al. 2019).

Hobday et al. (2018a) provided a refined definition that categorizes the severity of an MHW and allows for the identification of the more extreme events. Categories are delineated by the number of times the maximum observed temperature anomaly is greater than the difference between the climatological mean and the 90th-percentile threshold (**Figure 2d**). This approach results in a simple numbered system from category 1 (moderate) to 4 (extreme) that can be used to monitor an ongoing event or applied retrospectively given the maximum value reached.

Once a definition is chosen, a suite of metrics defines characteristics of each event. These metrics include intensity (e.g., temperature anomaly relative to the climatological mean), duration (time between start and end dates), and cumulative intensity (the intensity integrated over the duration of the event, analogous to degree heating days).

There are limitations with each type of MHW definition. The research questions and their application are important factors for selecting a definition, and the availability of long time series data is important for determining reference periods and thresholds.

3.2. Available Data Sets

Temperature data with daily resolution are ideal to identify and characterize MHWs in a consistent way. In addition, the data time period needs to be sufficiently long for estimating a climatology (ideally a minimum of 30 years; e.g., WMO 2018). However, Schlegel et al. (2019) showed that as few as 10 years of data may be sufficient for constraining the climatology, and recent studies have used fewer data, from short records or those with temporal gaps, to identify MHWs (e.g., Oliver et al. 2017).

Most studies have characterized MHWs using available SST products (see **Table 1**). Satellite-based SST data sets provide global, continuous, multidecadal, and near-real-time products for the study of MHWs. On local to regional scales, high-frequency subsurface ocean temperatures can be used to construct climatologies and analyze MHW characteristics (e.g., Schaeffer & Roughan 2017, Benthuisen et al. 2018, Elzahaby & Schaeffer 2019). However, subsurface measurements are generally available only in select locations (i.e., long-term moorings) or at a lower temporal resolution from depth-profiling drifting floats (e.g., Argo) or gliders. In fact, little is known about the characteristics of MHWs below the ocean surface due to lack of suitable data. Therefore, despite the notable importance of these events for benthic ecosystems, systematic studies of subsurface MHWs are difficult to undertake robustly.

When daily temperature data are not available (e.g., some seasonal forecasts and climate models save only monthly data), the ability to apply the formal MHW definitions is limited, given the necessity for daily measurements. However, such data have been used to examine longer-lasting MHWs in order to monitor coral bleaching risk (e.g., Donner et al. 2005) and perform seasonal forecasting of major events (Doi et al. 2013, Spillman et al. 2013, Jacox et al. 2019). In addition, the monthly temperature variations offer statistical insight into the frequency of extremes, a property that has been used to develop centennial-scale proxies of MHW properties globally (Oliver et al. 2018a).

In global ocean and climate models, SSTs and MHW estimates may have biases if subgrid-scale processes are not represented (e.g., Pilo et al. 2019) or do not include feedback from air–sea heat coupling (e.g., Frankignoul 1985). While data sets derived from observations, even coarsely resolved ones, inherently represent the physical processes of the natural world, coarse-resolution ocean models are challenged in representing ocean temperatures at the local scale. In global climate models, issues in representing boundary currents, eddies, teleconnections, coastal processes, and interannual-to-decadal variability will affect SST patterns (e.g., Cai & Cowan 2013, Taschetto et al. 2014, Sen Gupta et al. 2016, Power et al. 2017). These shortcomings may be mitigated by bias



Table 1 Ocean temperature data types along with their temporal coverage, spatial resolution, general strengths and weaknesses, and example products

Data set type	Earliest start date	Spatial resolution	Strengths	Weaknesses	Examples
Observations					
In situ (stationary)	Around 1900	Local	Accurate representation of local temperatures; several long records available; includes surface and subsurface measurements (in some cases)	Representation very spatially restricted; subject to missing data or changes in technology over time; some records too short to constrain a climatology	Scripps Shore Stations Program (USA), British Columbia Shore Station Oceanographic Program (Canada), Arendal (Norway), South African Coastal Temperature Network
In situ (passive movement)	Late twentieth century	Local to basin scale	Large spatial coverage; includes surface and subsurface measurements	Lagrangian in nature, which complicates analysis; instruments cannot be directed to sample a specific area of interest	Surface drifting buoys, Argo floats
In situ (active movement)	Early twenty-first century	Local to basin scale	Missions can target events as they occur; includes surface and subsurface measurements; other physical and biogeochemical parameters also typically measured	Data lack the spatiotemporal context outside the mission period and location; requires many missions to build up a time series for a particular location	Gliders, ships of opportunity (various missions and groups worldwide)
Remotely sensed (satellite: infrared, microwave)	1979	1–25 km	Global; continuous; near real time; high frequency (daily to weekly)	Records are too short for long-term climate change analyses; may not accurately resolve coastal temperatures	NOAA OI SST, CoralTemp, CCI-SST, GHRSSST
Statistically analyzed observations (bin averaged)	Late nineteenth century	5°	Global; long records; quantification of uncertainty	Low frequency (monthly); missing data in space and time	HadSST3
Statistically analyzed observations (interpolated)	Late nineteenth century	1–2°	Global; continuous; long records; quantification of uncertainty (in some cases)	Low frequency (monthly); data values in the absence of observations are weakly constrained by observations	HadISST, ERSST, COBE2
Reanalyses					
Eddy-resolving ocean reanalysis	1993	1/12–1/4°	Global; continuous; quantification of uncertainty (in some cases); complete three-dimensional ocean state estimated (temperature, salinity, and velocities)	Records are too short for long-term climate change analyses; data values in the absence of observations are weakly constrained by observations	GLORYS, HYCOM, BRAN

(Continued)



Table 1 (Continued)

Data set type	Earliest start date	Spatial resolution	Strengths	Weaknesses	Examples
Coarse-scale ocean reanalyses	Around 1900	1/2–1°	Global; continuous; long records; quantification of uncertainty (in some cases); complete three-dimensional ocean state estimated (temperature, salinity, and velocities)	Subgrid-scale physical processes are not resolved; data in the absence of observations, most notably in earlier time periods, are only very weakly constrained by observations	SODA, CERA-20C, GODAS
Numerical models					
Eddy-permitting and eddy-resolving ocean models	Various	1/4° or finer	Dynamically consistent data allow for an accurate exploration of physical processes; resolves important physical processes, including eddies and coastal processes	Free running models are not constrained to represent the timing of the observational record (i.e., are not useful for analyzing historical events)	Many modeling groups worldwide
Coarse-scale ocean models	Various	Coarser than 1/4°	Dynamically consistent data allow for an accurate exploration of physical processes	Free running models are not constrained to represent the timing of the observational record (i.e., are not useful for analyzing historical events); subgrid-scale physical processes are not resolved	Many modeling groups worldwide
Global and regional climate models	Various	1/2–2°	Dynamically consistent data allow for an accurate exploration of physical processes, including a representation of the complete, coupled climate system (ocean, atmosphere, cryosphere, etc.); can separate the natural and anthropogenic influences on MHWs	Free running models are not constrained to represent the timing of the observational record (i.e., are not useful for analyzing historical events); subgrid-scale physical processes are not resolved	CMIP5, CMIP6, CORDEX

Beggs (2020) provides further details on many specific observational SST data sets. Abbreviations: BRAN, Bluelink Reanalysis; CCI-SST, Climate Change Initiative Sea Surface Temperature; CERA-20C, Coupled Ocean-Atmosphere Reanalysis of the 20th Century; CMIP5/6, Coupled Model Intercomparison Project Phase 5/6; COBE2, Centennial In Situ Observation-Based Estimates 2; ERSST, Extended Reconstructed Sea Surface Temperature; GHRSSST, Group for High Resolution Sea Surface Temperature; GLORYS, Global Ocean Reanalysis; GODAS, Global Ocean Data Assimilation System; HadISST, Hadley Centre Sea Ice and Sea Surface Temperature; HadSST3, Hadley Centre Sea Surface Temperature; HYCOM, Hybrid Coordinate Ocean Model; MHW, marine heatwave; NOAA OI SST, National Oceanic and Atmospheric Administration Optimum Interpolation Sea Surface Temperature; SODA, Simple Ocean Data Assimilation; SST, sea surface temperature.

Surface mixed layer: the upper layer of the ocean, in contact with the atmosphere, which has a nearly uniform density over its depth due to turbulent mixing

Mixed-layer temperature budget: the theoretical balance of physical processes that cause temperature changes in the surface mixed layer

correction of the SST time series (e.g., Oliver et al. 2017, Pilo et al. 2019), but such an approach corrects only for the statistical aspects of the bias and not the underlying issues of poorly captured or unrepresented physical processes. Increases in model resolution and process parameterization may provide some improvements in this regard [e.g., Coupled Model Intercomparison Project Phase 6 (CMIP6); Eyring et al. 2016].

4. MARINE HEATWAVE MECHANISMS

4.1. Dynamical Understanding


The physical processes responsible for MHWs can be explored through the analysis of the heat sources and sinks within the surface mixed layer, which are reflected in SST variations (Alexander et al. 2000, Deser et al. 2010). A mixed-layer temperature budget has been used to describe the processes of MHW formation, evolution, and decay (Benthuisen et al. 2014, Chen et al. 2014, Kataoka et al. 2017). This approach relates temperature changes in the surface mixed layer with physical processes, including horizontal temperature transport and air–sea heat fluxes.

Generally, changes in mixed-layer temperatures arise from a combination of air–sea exchanges, advection by mean currents and eddies, horizontal and vertical mixing, and entrainment of water into the mixed layer. The rate of change of average seawater temperature in the mixed layer, or the temperature tendency, is given by (Moisan & Niiler 1998)

$$\begin{aligned}
 \underbrace{\frac{\partial \bar{T}}{\partial t}}_{\text{Temperature tendency}} = & - \underbrace{\bar{\mathbf{u}} \cdot \nabla \bar{T}}_{\text{Horizontal advection}} + \underbrace{\overline{\nabla \cdot (\kappa_h \nabla T)}}_{\text{Horizontal mixing}} - \underbrace{\frac{1}{b} \kappa_z \frac{\partial T}{\partial z} \Big|_{-b}}_{\text{Vertical mixing}} \\
 & - \underbrace{\left(\frac{\bar{T} - T_{-b}}{b} \right) \left(\underbrace{\frac{\partial b}{\partial t}}_{\text{MLD tendency}} + \underbrace{\bar{\mathbf{u}}_{-b} \cdot \nabla b}_{\text{Lateral induction}} + \underbrace{w_{-b}}_{\text{Vertical advection}} \right)}_{\text{Entrainment}} \\
 & + \underbrace{\frac{Q_{\text{SW}} - Q_{\text{SW}(-b)} + Q_{\text{LW}} + Q_{\text{sens}} + Q_{\text{lat}}}{\rho c_p b}}_{\text{Air-sea heat flux}}, \tag{1}
 \end{aligned}$$

where T is the temperature in the surface mixed layer, t is time, $\mathbf{u} = (u, v)$ is the two-dimensional horizontal (x, y) velocity vector, w is vertical (z) velocity, ∇ is the horizontal gradient operator, Q comprises various components of the air–sea heat flux (see details below), ρ is the seawater density, c_p is the specific heat capacity of seawater, b is the mixed-layer depth (MLD), and κ_h and κ_z are the horizontal and vertical diffusivity coefficients. Quantities have been vertically averaged over the mixed layer, and the vertical average of any quantity x is defined to be $\bar{x} = b^{-1} \int_{-b}^0 x dz$; a subscript x_{-b} indicates that the quantity is evaluated at the base of the mixed layer, i.e., at $z = -b$.

Equation 1 describes how the rate of change of surface mixed-layer temperature is related to the transfer of heat by horizontal advection, air–sea heat flux, lateral and vertical mixing, and entrainment of deeper waters into the mixed layer (**Figure 3a**). Horizontal temperature advection can drive local temperature changes through horizontal flows across a temperature gradient; examples include strong poleward geostrophic flows in a western boundary current extension region and strong Ekman flows associated with changes in wind stress (Rebert et al. 1985). Anomalous ocean currents—or, less often, anomalous temperature gradients—are responsible for




(Caption appears on following page)

Figure 3 (Figure appears on preceding page)

Physical processes affecting mixed-layer temperatures. (a) Relevant physical processes in a mixed-layer temperature budget. (b–g) Idealized examples of MHW events where the budget is dominated by (b–d) horizontal advection and (e–g) air–sea heat flux. For the advection event, we have (b) a background state consisting of a poleward ocean current \mathbf{u} (arrows) across a temperature gradient ∇T (colors; e.g., a western boundary current), which is (c) perturbed by an anomalous flow to the northeast, resulting in (d) anomalously high SSTs. For the air–sea heat flux event, we have (e) a background state consisting of an approximately stable stratification with a constant mixed-layer depth (b), wind forcing, and air–sea heat flux (Q). This state is (f) perturbed by an atmospheric event that acts to weaken the surface winds, increase the air–sea heat flux, and shoal the mixed-layer depth, resulting in (g) anomalously high temperatures in the mixed layer (and anomalously low temperatures immediately below the mixed layer, consistent with Sparnocchia et al. 2006). Abbreviations: MHW, marine heatwave; SST, sea surface temperature. Panel a adapted from Holbrook et al. (2019).

advective-type MHWs (**Figure 3b–d**), such as the 2015–2016 Tasman Sea MHW (Oliver et al. 2017) (**Figure 4a–g**). Vertical temperature advection results from vertical flows in the presence of thermal stratification and is often related to upwelling and downwelling processes. Schaeffer & Roughan (2017) showed subsurface coastal warming associated with downwelling-favorable winds, and Benthuisen et al. (2018) showed warm anomalies and subsurface MHW owing to anomalous downwelling via a reduction in the expected upwelling for that time of year. Upwelling

**Figure 4**

(Continued)




Figure 4

Anomalous contributions to the mixed-layer temperature budget during the (a–g) 2015–2016 Tasman Sea MHW and (h–n) 2003 Mediterranean Sea MHW. The maps show (a, b) SST anomaly, (b, i) mixed-layer depth (shading) and air–sea heat flux (Q) (contours, where red indicates a positive flux into the ocean) anomalies, and (c, j) velocity anomalies (arrows) and SST (shading) averaged over the peak duration of each event (December 1, 2015–February 29, 2016, for the Tasman Sea, and July 3–September 3, 2003, for the Mediterranean Sea). The time series show (d, k) SST anomaly, (e, l) Q anomaly, (f, m) mixed-layer depth anomaly, and (g, n) eastward and northward velocity anomalies spatially averaged. The time series are shaded or (in the case of velocities) the arrows colored according to the sign of the anomaly, with red and blue indicating a tendency to increase and decrease the mixed-layer temperature, respectively. Note that during the 2015–2016 Tasman Sea MHW, Q and the mixed-layer depth were variable but the velocities were predominantly southward, indicating that poleward advection of heat was a dominant contribution to the MHW. During the 2003 Mediterranean Sea MHW, the velocities were variable and weak, but the mixed-layer depths were anomalously shallow and Q tended to be positive, indicating a dominance of air–sea heat fluxes in the MHW development. Data are from GLORYS12V2 for SST, mixed-layer depth, and velocities and from ERA5 for Q ; anomalies are relative to the 1993–2013 climatology. Abbreviations: ERA5, European Centre for Medium-Range Weather Forecasts Reanalysis 5; GLORYS12V2, Global Ocean Reanalysis 12 version 2; MHW, marine heatwave; SST, sea surface temperature.

can also inhibit MHWs by bringing cool water to the surface (DeCastro et al. 2014, Gentemann et al. 2017, Fewings & Brown 2019).

The net air–sea heat fluxes (Q) are the sum of the net shortwave (Q_{SW}) and longwave (Q_{LW}) radiative components minus the fraction of shortwave radiation that escapes out the bottom of the mixed layer [$Q_{SW(-b)}$], as well as the latent (Q_{lat}) and sensible (Q_{sens}) turbulent heat fluxes (Cronin et al. 2019). With the exception of shortwave radiation, which has a depth structure, these heat

Serial autocorrelation: the correlation of a time series with a delayed version of itself

fluxes are distributed through the whole mixed layer, and so their effect on temperature tendency is inversely related to the mixed-layer depth (b). MHWs are often associated with anomalous air–sea heat fluxes, which can include abnormally high Q_{SW} , as a result of less cloud cover and greater insolation, or Q_{sens} when the surface air is warm. Often, both scenarios occur during an atmospheric high-pressure system and coincide with reduced wind speeds that suppress vertical mixing, thereby reducing b . Meanwhile, unusually low latent heat loss from the ocean (negative Q_{lat}), as a result of weak winds, may also contribute to temperature increases. These processes, which may act independently or simultaneously, are responsible for air–sea heat flux–type MHWs (Figure 3e,f), such as the 2003 Mediterranean Sea MHW (Sparnocchia et al. 2006, Olita et al. 2007) (Figure 4b–n).

The remaining terms in Equation 1 are associated with mixing due to horizontal diffusive flux, mixing due to vertical turbulent flux at the base of the mixed layer, and entrainment of waters into the mixed layer due to temporal or spatial variations in mixed-layer depth. These processes are often assumed to account for a smaller proportion of mixed-layer temperature changes associated with MHWs, and so are often neglected or considered part of a residual term.

4.2. Statistical Understanding

The statistical properties of a daily temperature time series form the basis of MHW properties, such as their frequency, duration, and intensity. We can decompose the time series of temperature as

$$T_t = T_t^{\text{tr}} + T_t^{\text{S}} + T_t^{\text{NS}}, \quad 2.$$

where T_t is the SST at time t , T_t^{tr} is the change related to the long-term (secular) trend, T_t^{S} is the seasonal climatological mean (identically repeating each year), and T_t^{NS} is the nonseasonal component of the SST (the anomalies, with the secular trend removed). If MHWs are defined as SST values above a seasonally varying threshold (e.g., Hobday et al. 2016), and assuming that there is no long-term trend in SST, then MHW SSTs are dependent on the nonseasonal component's properties.

Frankignoul & Hasselmann (1977) developed a stochastic climate model to represent the temperature of a one-dimensional surface mixed layer forced by noisy surface heat fluxes. This model elucidates the relationships between MHW properties and the statistical characteristics of SST. Considering the nonseasonal component T_t^{NS} to have zero mean, variance σ^2 , and a nonnegligible serial autocorrelation (i.e., the memory timescale τ), the nonseasonal component can be modeled as an order-1 autoregressive (AR1) process (see also Deser et al. 2010), so that the temperature time series is red noise:

$$T_{t+1}^{\text{NS}} = aT_t^{\text{NS}} + \epsilon_t, \quad 3.$$

where a is the autoregressive parameter ($0 \leq a \leq 1$), and ϵ_t is a white noise process that is assumed here to be normally distributed with zero mean and variance σ_ϵ^2 . The memory timescale of the AR1 process τ increases with a , given by $\tau = -1/\ln a$, and the variance of T_t^{NS} is a function of both a and σ_ϵ^2 , given by $\sigma^2 = \sigma_\epsilon^2/(1 - a^2)$. Given values of a and σ_ϵ^2 and realizations of ϵ_t , simulations of T_t^{NS} and (after assuming a sinusoidal form for T_t^{S}) T_t may be generated (Figure 5a–c).

Memory timescales are related to the ocean's slow integration of relatively fast atmospheric forcing (e.g., atmospheric weather on timescales of one to two weeks) via surface heat fluxes, near-surface currents, and mixing processes, and temperatures relax to climatological values via damping caused by turbulent energy and longwave radiative fluxes (Deser et al. 2010, Di Lorenzo & Ohman 2013). The variance of this process is related to energetics of the atmospheric forcing and




Figure 5

The effect of SST statistical properties on MHW characteristics. (a–c) Simulated SST time series assuming a sinusoidal seasonal climatology and a nonseasonal component given by an AR1 process with memory timescale τ and noise variance $\sigma_\epsilon^2 = (1 - a^2)\sigma^2$. Shaded areas indicate MHWs according to the Hobday et al. (2016) definition. (d–f) MHW properties as a function of SST standard deviation σ and memory timescale τ . The plots show the annual maximum MHW intensity (panel d), annual mean MHW duration (panel e), and annual count of MHW events (panel f). An ensemble of 100 SST time series were simulated for each σ_ϵ and τ value over the ranges 0.01–2.00°C and 10–100 days, respectively. Abbreviations: AR1, order-1 autoregressive; MHW, marine heatwave; SST, sea surface temperature.

Stationarity:

the condition that the statistical properties of a time series (e.g., mean and variance) do not change over time

the local ocean dynamics (e.g., the generation of eddies via instability growth). MHWs occurring in time series with larger τ (Figure 5c) are typically longer than for those from time series with smaller τ (Figure 5a,b). In addition, MHWs are more intense for larger σ^2 (Figure 5b,c) than for smaller σ^2 (Figure 5a). Performing simulations of an ensemble of 40-year SST time series for a range of realistic values of σ^2 and τ and calculating the ensemble-mean annual MHW properties shows that these properties are strongly dependent on underlying SST properties. Specifically, annual maximum MHW intensity increases with SST variance (Figure 5d), annual mean MHW duration increases with SST memory timescale (Figure 5e), and annual MHW count (frequency) decreases with SST memory timescale (Figure 5f).

The conclusions provided above assume that the SST statistics are stationary, i.e., that the mean and variance of SST do not change over time. In a warming ocean, we know that this is unlikely to be true, particularly for the mean SST (Oliver 2019). Therefore, we expect trends in the mean SST (effectively a nonzero T_t'') and SST variance to affect MHW properties. Nonstationarity in SST statistics leads directly to trends in MHW frequency, duration, and intensity (Oliver et al. 2018a, Oliver 2019). This aspect is an important consideration that must be taken into account when defining baseline periods for studies on the effects of climate change on MHWs (see also Sections 5.3, 5.4, and 6.1).

5. MEAN STATE, VARIABILITY, AND LONG-TERM TRENDS IN MARINE HEATWAVES

5.1. Global Distribution

Remotely sensed SSTs indicate that MHW frequencies range from approximately one to three events per year on average (Figure 6a). In the eastern tropical Pacific, however, El Niño–Southern Oscillation (ENSO) events manifest as individual, long-lasting MHWs (Holbrook et al. 2020a). Hot spots of high MHW intensity (Figure 6b) occur in regions of large SST variability (Figure 6e), including the five western boundary current extension regions (Chen et al. 2014, Oliver et al. 2017), the central and eastern equatorial Pacific Ocean (Echevin et al. 2018), and eastern boundary current regions (Rouault et al. 2007). Typical MHW durations (Figure 6c) are longest in the eastern tropical Pacific, a region dominated by ENSO SST variability with an average duration of up to 60 days (Holbrook et al. 2020a), and shortest over other tropical regions, typically 5–10 days. In the extratropics, MHW durations are more uniformly 10–15 days, with the northeast and southeast Pacific Ocean being exceptions (up to 30-day mean durations; Di Lorenzo & Mantua 2016).

The observed spatial pattern of MHW properties is dependent on the physical processes controlling temperature variability across the globe (see Section 4.1). Unstable boundary currents and mesoscale eddies cause MHWs through anomalous temperature advection in those regions, and anomalous air–sea heat flux is responsible for setting MHW properties in regions susceptible to atmospheric forcing. The observed patterns of SST and MHW statistics follow the expected statistical relationships (see Section 4.2). Spatial correlations show that MHW intensity is strongly correlated with SST variance ($r = 0.95$) and MHW duration is strongly related to SST memory timescale ($r = 0.68$).

5.2. Variability

MHWs are modulated by local and remote processes acting across a large range of spatial and temporal scales (see figure 2 in Holbrook et al. 2019). The strength of many of these local processes, including surface heat fluxes and vertical mixing, are a function of the overlying atmospheric synoptic conditions (winds, cloud cover, and humidity). These conditions may in






Figure 6

Statistical properties of (a–c) MHWs and (d–f) SST. The plots shown are the (a) annual count of MHW days, (b) annual maximum MHW intensity, (c) annual mean MHW duration, (d) annual mean SST, (e) standard deviation of the nonseasonal component of SST, and (f) memory timescale of the nonseasonal component of SST (Section 4.2). The nonseasonal component of SST was calculated by subtracting the seasonal climatology and long-term linear trend. The plots were created using NOAA OI SST V2.0 data for 1982–2019, with a baseline period of 1982–2011. Abbreviations: MHW, marine heatwave; NOAA OI SST V2.0, National Oceanic and Atmospheric Administration Optimum Interpolation Sea Surface Temperature version 2.0; SST, sea surface temperature.

turn vary with changes in the general circulation (e.g., the Walker circulation) or planetary waves generated remotely (e.g., associated with ENSO or the Madden–Julian Oscillation). Such remotely forced variations can cause oceanic disturbances that propagate large distances (via Rossby and Kelvin waves), affecting local ocean advection of temperature with long predictability timescales (Holbrook et al. 2019, 2020b).

ENSO is the dominant mode of interannual climate variability across the globe (e.g., McPhaden et al. 2006). It is also a leading cause of MHW occurrences globally (Oliver et al. 2018a), with key influences and impacts in the Indo-Pacific (Holbrook et al. 2019, 2020a) and more distant links to the Atlantic and Southern Oceans (Holbrook et al. 2019). Coupled air–sea interactions in the tropical Pacific are related to large shifts in SST and directly affect MHW

Anthropogenic forcing: forcing acting on the Earth's climate system that is caused by human activities (e.g., greenhouse gas emissions, aerosol emissions, and land use changes)

Natural variability: spatiotemporal variations in climate that are caused by natural forcing (e.g., solar, volcanic, and orbital forcing) acting on the Earth's climate system

characteristics in the central and eastern tropical Pacific. ENSO teleconnections via atmospheric and oceanic pathways have been an important factor in triggering major MHWs in subtropical to midlatitude regions of the northeast Pacific (e.g., Doi et al. 2015; Di Lorenzo & Mantua 2016; Jacox et al. 2017, 2019), in the central South Pacific (Lee et al. 2010), and in the eastern Indian Ocean off Australia's west coast (Doi et al. 2013, Feng et al. 2013, Benthuyssen et al. 2014, Marshall et al. 2015). Such ENSO teleconnections can be amplified by local air–sea feedback processes (e.g., Marshall et al. 2015, Myers et al. 2018) or work in concert with stochastic local weather conditions.

Other climate modes also modulate the occurrence rate of MHWs either locally where the mode operates or in remote regions via teleconnections (Holbrook et al. 2019). Significant relationships exist among indices of the Indian Ocean Dipole (the Dipole Mode Index), ENSO (the El Niño Modoki and Niño3.4 indices; see figures 3*b* and 4 in Holbrook et al. 2019), and MHW occurrences and likelihoods across the tropical Indo-Pacific region. These modes exhibit a degree of predictive skill, suggesting that it may be possible to make MHW probabilistic forecasts (Holbrook et al. 2020*b*) using knowledge of ENSO flavors (e.g., Capotondi & Sardeshmukh 2015, Capotondi et al. 2015), the Indian Ocean Dipole (e.g., Zhao et al. 2019), and the Madden–Julian Oscillation (Zhang et al. 2017, Lim et al. 2018).

5.3. Long-Term Trends

Given the dependence of MHW characteristics on the underlying SST properties, along with a warming ocean, significant changes in MHW characteristics have been identified over the historical record. Notably, increases in both the mean SST and the variability of SST can lead to increases in warm temperature extremes (Field et al. 2012). Much of the global ocean has seen mean SST warming over the satellite record, and several such hot spots have also seen increases in SST variability, with implications for changes in MHWs (**Figure 7**).


The days of extremely high SST have become more common over 38% of the world's coastline (Lima & Wetthey 2012). As a global average, the frequency, intensity, duration, and spatial extent of MHWs have increased substantially over the satellite record (Frölicher et al. 2018, Oliver et al. 2018*a*). MHW frequency increased over 82% of the global ocean but decreased in parts of the Southern Ocean (Oliver et al. 2018*a*). The intensity of MHWs has increased most notably in the five midlatitude western boundary current extension regions, areas noted to be warming considerably faster than the global mean rate (Wu et al. 2012, Hu et al. 2015). The global-average duration of MHWs has nearly doubled over the satellite record, particularly over the mid- and high-latitude regions of all ocean basins (Oliver et al. 2018*a*).

While a few studies have assessed longer-term changes in MHWs over the presatellite records, these analyses are limited in overall spatiotemporal coverage and homogeneity by data availability (see Section 3.2). Using monthly SSTs as a proxy for MHW properties, Oliver et al. (2018*a*) showed that the annual count of MHW days has increased by more than 50% since the early twentieth century. Several coastal locations around the world have time series of daily or weekly SSTs (see **Table 1**), which could provide additional estimates of long-term changes in coastal MHWs.

5.4. The Role of Anthropogenic Climate Change

Long-term trends in MHW characteristics (frequency, intensity, and duration) over the instrumental record are strongly linked to anthropogenic climate change. In global climate model simulations, substantial increases in global MHW frequency, intensity, duration, and spatial extent were consistent with anthropogenic forcing (Frölicher et al. 2018), with properties exceeding the range expected by natural variability in the early-to-mid-twenty-first century (Oliver et al. 2019).

Changes in global MHW characteristics have been strongly linked to changes in the background SST (Oliver 2019); these increases are expected to continue under projected future emissions scenarios, and it is possible that much of the global ocean will reach a permanent MHW state by the late twenty-first century (Oliver et al. 2019). Most studies on the role of climate change have examined MHWs globally, but there is a need to examine regional cases. Such analyses have been



(Caption appears on following page)

Figure 7 (Figure appears on preceding page)

Change in SST statistics and relationship to changes in extreme temperatures. The map shows where SSTs indicate an increase in annual mean SST (*light red*) and SST variance (*dark red*) over the satellite period. White indicates no increase in either; an increase is measured by a positive linear trend in NOAA OI SST V2.0 over 1982–2017. The four surrounding panels show probability density functions of daily SST anomalies from four regions (matching four of the regions shown in **Figure 1a**). The functions shown are for the full time series (*dashed gray lines*), the first half of the data (*solid black lines*), and the second half of the data (*solid red lines*); shaded regions indicate the area of the function above the 90th percentile of the full time series. The mean (μ , in degrees Celsius) and standard deviation (σ , in degrees Celsius) are also shown for the two time periods. Abbreviations: NOAA OI SST V2.0, National Oceanic and Atmospheric Administration Optimum Interpolation Sea Surface Temperature version 2.0; SST, sea surface temperature.

performed for a few regions, including the North Atlantic (Alexander et al. 2018) and the Mediterranean Sea (Darmaraki et al. 2019), but such analyses are encouraged for other regions as well.

A growing body of literature has examined how individual MHW events are attributable to global warming (see also Collins et al. 2019). These analyses draw from a method of event attribution, based on the fraction of attributable risk, that is used in atmospheric heatwave studies (Stott et al. 2004). To date, event attribution studies have been performed on the 2015–2016 Tasman Sea event (Oliver et al. 2017), the 2017–2018 Tasman Sea event (Perkins-Kirkpatrick et al. 2019), the 2016 event across northern Australian waters (Oliver et al. 2017), the 2016 event in the Gulf of Alaska and Bering Sea (Oliver et al. 2018b, Walsh et al. 2018), the northeast Pacific Blob (Weller et al. 2015), the 2015–2016 extreme El Niño warming in the central equatorial Pacific (Newman et al. 2018), and the 2016 California Current event (Jacox et al. 2017). Some studies have extended attribution approaches to include the likelihood of ecosystem impacts, such as Great Barrier Reef coral bleaching in 2016 (Lewis & Mallela 2018). In general, anthropogenic climate change has substantially increased the likelihood of MHWs with the observed intensity and duration, although several notable MHWs remain to be attributed (Collins et al. 2019). Most studies attribute the temperature anomalies associated with the events (the intensity), but a few also independently attribute the duration of the event. Presently, there is no consistent framework for jointly attributing the totality of the event characteristics.

6. CONTEMPORARY ISSUES

6.1. Baseline Periods for Marine Heatwave Analyses

Given significant ocean warming, a major issue for MHW analyses is choosing the baseline period for calculating the climatological mean and percentile metrics. A fixed-baseline period is commonly used in climatology studies (WMO 2017, 2018) and is standard practice in atmospheric studies (Perkins et al. 2012, Perkins & Alexander 2013). Many MHW studies have chosen fixed-baseline periods (e.g., Liu et al. 2014, Benthuyssen et al. 2018, Oliver et al. 2018a), while others have recommended a moving baseline (Jacox 2019). The baseline period affects how the long-term trend in mean SST is expressed through MHWs (Section 4.2), and specific research questions of interest dictate the decision to use a fixed versus moving baseline.

The effect of a fixed versus moving baseline period can be demonstrated using a time series of SST exhibiting a long-term trend. To do so, we used SST area-averaged over a region in the northeast Pacific from the Community Earth System Model (CESM) Large Ensemble Numerical Simulation (LENS) climate model (Kay et al. 2015) historical and Representative Concentration Pathway 8.5 future projection experiments, which together span 1920–2080 (**Figure 8a**). We used two different approaches to calculate the SST climatology and anomaly time series: one using a fixed 31-year baseline period and one using a moving 31-year baseline centered on the year in question (**Figure 8b,c**). We then used the exceedances above the resulting thresholds (**Figure 8c**) to determine MHWs following the Hobday et al. (2016) method. A fixed-baseline period leads

Event attribution:

a quantification of the role of anthropogenic versus natural climate forcing in the probability of extreme climate events, often with the use of global climate model experiments




Figure 8

The effect of using a fixed baseline versus a moving baseline on the calculation of SST climatologies, anomalies, and MHWs. The plots shown are time series of (a) SST, (b) SST anomalies, (c) the climatological 90th-percentile threshold, and (d) annual MHW days; panels b–d were calculated using either a fixed baseline (1980–2010; blue lines) or a moving contemporary baseline (red lines). The initial time series in panel a is SST area-averaged over 30–57°N, 140–120°W (the Blob region), from the historical and Representative Concentration Pathway 8.5 simulations of the CESM LENS global climate model (1920–2080). Abbreviations: CESM, Community Earth System Model; LENS, Large Ensemble Numerical Simulation; MHW, marine heatwave; SST, sea surface temperature.

to a saturation of SST above the threshold, which in turn leads to a full year of MHW days (a permanent MHW state) in the later part of the twenty-first century, while a moving-baseline period leads to more stationary MHW properties over time (**Figure 8d**). The use of a fixed-baseline period assigns all of the long-term warming to the anomalies and thus to the MHWs (i.e., in the notation of Section 4.2, the anomalies represent $T_t^w + T_t^{NS}$ and the climatology only T_t^S), while the use of a moving baseline assigns all of the long-term warming to the climatology (and thus removes it from the anomalies and MHWs—i.e., the anomalies represent only T_t^{NS} and the climatology $T_t^S + T_t^w$).

Both methods of calculating baseline climatologies are complementary, and the recommended choice depends on the questions asked. Much of the study of MHWs is motivated by ecological impacts, which raises the issue of each species' rate of adaptation. The rate and degree of ecosystem change under warming conditions depend on the thermal tolerance ranges and diversity of particular species (Stuart-Smith et al. 2015, Burrows et al. 2019). However, major questions remain as to whether species can acclimatize and adapt to warmer conditions and, if so, how quickly they can

do so (Donelson et al. 2019, Fox et al. 2019). A fixed-baseline period is particularly useful when studying ecosystems with a slow (or no) capacity for adaptation and for illustrating overall changes, including long-term warming effects. Moving-baseline periods are most useful when considering ecosystems with a faster adaptation capacity or greater mobility (e.g., the ability to move to new habitats) and for studying physical climate characteristics with a focus on variability rather than long-term warming, as this approach primarily reflects shorter-term variability changes (changes in temperature variance and skewness). However, further research on species' and ecosystems' acclimatization and adaptation timescales are needed to inform these choices. The combination of both approaches (i.e., considering both in parallel) can be useful to disentangle the effects of long-term warming from changes in the magnitude of variability.

6.2. Communicating About Marine Heatwaves

A contemporary issue in many branches of science is effective communication. The science of understanding MHWs will be advanced with a clear communication approach to the description of events, which will aid public communication, allied research communities, and funding initiatives. As with other extreme events, including atmospheric heatwaves, tropical cyclones, and bushfires, naming and categorizing MHWs allows for straightforward communication (Hobday et al. 2018a). Importantly, naming an event based on the year and location ensures that extreme events are consistently discussed. A categorization system based on an MHW's maximum intensity allows for monitoring and communicating how its characteristics are changing in near real time.

Real-time tracking and monitoring of MHWs also offer a communication opportunity. The rapid deployment of ocean gliders during ongoing events and the dissemination of near-real-time data are expected to be of considerable interest and potential value to marine managers and the tourism, fishing, and aquaculture industries (for several examples of MHW tracking tools, see the Related Resources section at the end of this article).

There is demand for information about future weather and climate, including subseasonal to seasonal forecasts of MHWs. These forecasts are on timescales that allow proactive planning rather than reactive responses (Hobday et al. 2018b) and are based on statistical relationships and dynamical models (Holbrook et al. 2020b) (for examples, see the Related Resources section). Recent developments are enhancing understanding of the physical mechanisms that give rise to MHWs, which underpin prediction systems of MHWs (Jacox et al. 2020). Using climate mode indices may enable successful climate forecasts of likely increases or decreases in MHW days over broad geographic areas (Holbrook et al. 2019).

Changes in extreme events are strong expressions of anthropogenic climate change. Given the relatively small global-average, multidecadal-to-centennial-timescale temperature rise that climate change represents (e.g., the Paris Agreement target of limiting the total increase to $<2^{\circ}\text{C}$), the public has a mixed awareness of the impacts of climate change on temperature [i.e., across the full temperature probability distribution, including the extremes (tails)] and the associated climate feedbacks and ecosystem impacts. It is clear that MHWs, as extreme events with dramatic effects, can help scientists and policy makers to build public understanding about the urgency of response to long-term environmental change.

SUMMARY POINTS

1. There are a diversity of approaches for how to define marine heatwaves (MHWs), using fixed, relative, or seasonally varying thresholds, and each approach has its advantages

and disadvantages. Consistent definitions and baselines facilitate comparisons and will accelerate learning about MHWs in different parts of the world.

2. Global observations of sea surface temperature by satellite and advances in data assimilative ocean models have made it feasible to monitor MHWs on a global scale in near real time.
3. The physical drivers of MHWs can be elucidated using a mixed-layer temperature budget, which relates the accumulation of surface ocean heat to physical processes, including horizontal currents and air–sea heat fluxes.
4. MHW characteristics are linked to the statistical properties of the temperature time series, with event intensity increasing with temperature variance and event duration increasing with serial autocorrelation.
5. MHWs are distributed globally and have different characteristics depending on the regionally dominant physical processes and the statistics of the underlying temperature distribution. For example, highly variable regions, such as western boundary extensions, have high-intensity events that are caused predominantly by anomalous heat transport.
6. The historical record shows that the occurrence frequency of MHWs and their characteristics are sensitive to climate modes of variability and long-term warming, which has implications for the predictability of MHW events.
7. Anthropogenic climate change has led to substantial increases in the intensity and duration of MHWs, with several recent events explicitly attributable to anthropogenic forcing, and this is projected to continue throughout the twenty-first century.

FUTURE ISSUES

1. Efforts toward developing a process-based understanding of MHW mechanisms and their predictability need to inform the development of appropriate forecast systems with relevant lead times.
2. Reliable, consistent observational data sets of daily ocean temperatures, including subsurface temperatures, are needed to study observed MHWs with improved data coverage in space and time.
3. MHW properties have been studied based on available data primarily at the surface, and understanding of their manifestation at depth is more limited. Further research is needed to develop a more robust understanding of the physical processes and climate drivers of subsurface MHWs and their links to the surface.
4. To better inform climatological baseline periods, further research is required on linking the physical and climatological aspects of MHWs with species' and ecosystems' acclimatization and adaptation timescales.
5. In the case of MHWs calculated with a fixed baseline and in the presence of long-term warming conditions, is the notion of permanent MHW conditions useful? How can ecosystem adaptation timescales be incorporated into how baselines are defined?



6. In regions where the environmental conditions have radically and abruptly shifted (e.g., in areas historically dominated by sea ice cover), there is no clear definition of MHWs.
7. Currently, global systems for SST and other data streams are in place that allow for near-real-time identification of MHWs. These systems can be adapted to ensure that when unprecedented MHWs arise, resources are available to the public and the scientific community for rapid research and dissemination of information regarding ongoing events.

DISCLOSURE STATEMENT

The authors are not aware of any affiliations, memberships, funding, or financial holdings that might be perceived as affecting the objectivity of this review.

ACKNOWLEDGMENTS

Through the years, our understanding of MHWs has evolved through valuable discussions and/or collaborations with a broad group of people, including, but not limited to, Lisa Alexander, Nathan Bindoff, Michael Burrows, Madeleine Cahill, Nick Caputi, Ke Chen, Ming Feng, Michael Jacox, Scott Ling, Andrew Marshall, Martin Marzloff, Pippa Moore, Craig Mundy, Sarah Perkins-Kirkpatrick, Hillary Scannell, Dan Smale, Sandra Straub, Mads Thomsen, Thomas Wernberg, and Simon Wotherspoon. In particular, we would like to recognize the Marine Heatwaves International Working Group (<http://www.marineheatwaves.org>) and several of its workshops for fostering much of this discussion and research.

E.C.J.O. received support for this work from National Sciences and Engineering Research Council of Canada Discovery Grant RGPIN-2018-05255, the Ocean Frontier Institute through an award from the Canada First Research Excellence Fund, and Marine Environmental Observation, Prediction, and Response Network Early Career Faculty Grant 1-02-02-059.1. M.G.D. receives support from Spanish Ministry for the Economy, Industry, and Competitiveness grant reference RYC-2017-22964. N.J.H. acknowledges support from the Australian Research Council Centre of Excellence for Climate Extremes and the Australian government's National Environmental Science Programme Earth Systems and Climate Change Hub.

LITERATURE CITED

- Alexander MA, Scott JD, Deser C. 2000. Processes that influence sea surface temperature and ocean mixed layer depth variability in a coupled model. *J. Geophys. Res. Oceans* 105:16823–42
- Alexander MA, Scott JD, Friedland KD, Mills KE, Nye JA, et al. 2018. Projected sea surface temperatures over the 21st century: changes in the mean, variability and extremes for large marine ecosystem regions of northern oceans. *Elem. Sci. Anthr.* 6:9
- Beggs H. 2020. Temperature. In *Earth Observation: Data, Processing and Applications*, Vol. 3B: *Applications—Surface Waters*, ed. BA Harrison, JA Anstee, A Dekker, S Phinn, N Mueller, G Byrne. Melbourne, Aust.: Coop. Res. Cent. Spat. Inf. In press
- Benthuisen JA, Feng M, Zhong L. 2014. Spatial patterns of warming off Western Australia during the 2011 Ningaloo Niño: quantifying impacts of remote and local forcing. *Cont. Shelf Res.* 91:232–46
- Benthuisen JA, Oliver ECJ, Chen K, Wernberg T. 2020. Editorial: advances in understanding marine heatwaves and their impacts. *Front. Mar. Sci.* 7:147
- Benthuisen JA, Oliver ECJ, Feng M, Marshall AG. 2018. Extreme marine warming across tropical Australia during austral summer 2015–2016. *J. Geophys. Res. Oceans* 123:1301–26

- Bond NA, Cronin MF, Freeland H, Mantua N. 2015. Causes and impacts of the 2014 warm anomaly in the NE Pacific. *Geophys. Res. Lett.* 42:3414–20
- Burrows MT, Bates AE, Costello MJ, Edwards M, Edgar GJ, et al. 2019. Ocean community warming responses explained by thermal affinities and temperature gradients. *Nat. Clim. Change* 9:959–63
- Cai W, Cowan T. 2013. Why is the amplitude of the Indian Ocean Dipole overly large in CMIP3 and CMIP5 climate models? *Geophys. Res. Lett.* 40:1200–5
- Capotondi A, Sardeshmukh PD. 2015. Optimal precursors of different types of ENSO events. *Geophys. Res. Lett.* 42:9952–60
- Capotondi A, Wittenberg A, Newman M, Di Lorenzo E, Yu J, et al. 2015. Understanding ENSO diversity. *Bull. Am. Meteorol. Soc.* 96:921–38
- Caputi N, Kangas M, Denham A, Feng M, Pearce A, et al. 2016. Management adaptation of invertebrate fisheries to an extreme marine heat wave event at a global warming hot spot. *Ecol. Evol.* 6:3583–93
- Cavanaugh KC, Dangremond EM, Doughty CL, Williams AP, Parker JD, et al. 2019. Climate-driven regime shifts in a mangrove–salt marsh ecotone over the past 250 years. *PNAS* 116:21602–8
- Chen K, Gawarkiewicz GG, Kwon Y-O, Zhang WG. 2015. The role of atmospheric forcing versus ocean advection during the extreme warming of the northeast U.S. continental shelf in 2012. *J. Geophys. Res. Oceans* 120:4324–39
- Chen K, Gawarkiewicz GG, Lentz SJ, Bane JM. 2014. Diagnosing the warming of the northeastern U.S. coastal ocean in 2012: a linkage between the atmospheric jet stream variability and ocean response. *J. Geophys. Res. Oceans* 119:218–27
- Coles S, Bawa J, Trenner L, Dorazio P. 2001. *An Introduction to Statistical Modeling of Extreme Values*. London: Springer
- Collins M, Sutherland M, Bouwer L, Cheong S-M, Frölicher T, et al. 2012. Extremes, abrupt changes and managing risks. In *IPCC Special Report on the Ocean and Cryosphere in a Changing Climate*, ed. H-O Pörtner, DC Roberts, V Masson-Delmotte, P Zhai, M Tignor, et al., pp. 589–655. Geneva: Intergov. Panel Clim. Change
- Cronin MF, Gentemann CL, Edson JB, Ueki I, Bourassa M, et al. 2019. Air-sea fluxes with a focus on heat and momentum. *Front. Mar. Sci.* 6:430
- Darmaraki S, Somot S, Sevault F, Nabat P, Narvaez WDC, et al. 2019. Future evolution of marine heatwaves in the Mediterranean Sea. *Clim. Dyn.* 53:1371–92
- DeCastro M, Gómez-Gesteira M, Costoya X, Santos F. 2014. Upwelling influence on the number of extreme hot SST days along the Canary upwelling ecosystem. *J. Geophys. Res. Oceans* 119:3029–40
- Deser C, Alexander MA, Xie S-P, Phillips AS. 2010. Sea surface temperature variability: patterns and mechanisms. *Annu. Rev. Mar. Sci.* 2:115–43
- Di Lorenzo E, Mantua N. 2016. Multi-year persistence of the 2014/15 North Pacific marine heatwave. *Nat. Clim. Change* 6:1042–47
- Di Lorenzo E, Ohman MD. 2013. A double-integration hypothesis to explain ocean ecosystem response to climate forcing. *PNAS* 110:2496–99
- Doi T, Behera S, Yamagata T. 2013. Predictability of the Ningaloo Niño/Niña. *Sci. Rep.* 3:2892
- Doi T, Yuan C, Behera SK, Yamagata T. 2015. Predictability of the California Niño/Niña. *J. Clim.* 28:7237–49
- Donelson JM, Sunday JM, Figueira WF, Gaitán-Espitia JD, Hobday AJ, et al. 2019. Understanding interactions between plasticity, adaptation and range shifts in response to marine environmental change. *Philos. Trans. R. Soc. B* 374:20180186
- Donner SD, Skirving WJ, Little CM, Oppenheimer M, Hoegh-Guldberg O. 2005. Global assessment of coral bleaching and required rates of adaptation under climate change. *Glob. Change Biol.* 11:2251–65
- Eakin CM, Morgan JA, Heron SF, Smith TB, Liu G, et al. 2010. Caribbean corals in crisis: record thermal stress, bleaching, and mortality in 2005. *PLOS ONE* 5:e13969
- Eakin CM, Sweatman HP, Brainard RE. 2019. The 2014–2017 global-scale coral bleaching event: insights and impacts. *Coral Reefs* 38:539–45
- Echevin V, Colas F, Espinoza-Morriberon D, Vasquez L, Anculle T, Gutierrez D. 2018. Forcings and evolution of the 2017 coastal El Niño off Northern Peru and Ecuador. *Front. Mar. Sci.* 5:367



- Elzahaby Y, Schaeffer A. 2019. Observational insight into the subsurface anomalies of marine heatwaves. *Front. Mar. Sci.* 6:745
- Eyring V, Bony S, Meehl GA, Senior CA, Stouffer RJ, Taylor KE. 2016. Overview of the Coupled Model Inter-comparison Project Phase 6 (CMIP6) experimental design and organization. *Geosci. Model Dev.* 9:1937–58
- Feng M, McPhaden M, Xie S, Hafner J. 2013. La Niña forces unprecedented Leeuwin Current warming in 2011. *Sci. Rep.* 3:1277
- Fewings MR, Brown KS. 2019. Regional structure in the marine heat wave of summer 2015 off the western United States. *Front. Mar. Sci.* 6:564
- Field CB, Barros V, Stocker TF, Dahe Q, Dokken DJ, eds. 2012. *Managing the Risks of Extreme Events and Disasters to Advance Climate Change Adaptation: Special Report of the Intergovernmental Panel on Climate Change*. Cambridge, UK: Cambridge Univ. Press
- Firth LB, Knights AM, Bell SS. 2011. Air temperature and winter mortality: implications for the persistence of the invasive mussel, *Perna viridis* in the intertidal zone of the south-eastern United States. *J. Exp. Mar. Biol. Ecol.* 400:250–56
- Fordyce AJ, Ainsworth TD, Heron SF, Leggat W. 2019. Marine heatwave hotspots in coral reef environments: physical drivers, ecophysiological outcomes and impact upon structural complexity. *Front. Mar. Sci.* 6:498
- Fox RJ, Donelson JM, Schunter C, Ravasi T, Gaitán-Espitia JD. 2019. Beyond buying time: the role of plasticity in phenotypic adaptation to rapid environmental change. *Philos. Trans. R. Soc. B* 374:20180174
- Frankignoul C. 1985. Sea surface temperature anomalies, planetary waves, and air-sea feedback in the middle latitudes. *Rev. Geophys.* 23:357–90
- Frankignoul C, Hasselmann K. 1977. Stochastic climate models, part II: application to sea-surface temperature anomalies and thermocline variability. *Tellus* 29:289–305
- Frölicher TL, Fischer EM, Gruber N. 2018. Marine heatwaves under global warming. *Nature* 560:360–64
- Frölicher TL, Laufkötter C. 2018. Emerging risks from marine heat waves. *Nat. Commun.* 9:650
- Garrabou J, Coma R, Bensoussan N, Bally M, Chevaldonné P, et al. 2009. Mass mortality in Northwestern Mediterranean rocky benthic communities: effects of the 2003 heat wave. *Glob. Change Biol.* 15:1090–103
- Gawarkiewicz G, Chen K, Forsyth J, Bahr F, Mercer AM, et al. 2019. An advective marine heatwave in the Middle Atlantic Bight: shelfbreak exchange driven thermohaline anomalies in early 2017. *Front. Mar. Sci.* 6:712
- Gentemann CL, Fewings MR, Garca-Reyes M. 2017. Satellite sea surface temperatures along the West Coast of the United States during the 2014–2016 northeast Pacific marine heat wave. *Geophys. Res. Lett.* 44:312–19
- Gumbel EJ. 1958. *Statistics of Extremes*. New York: Columbia Univ. Press
- Gunter G. 1951. Destruction of fishes and other organisms on the south Texas coast by the cold wave of January 28–February 3, 1951. *Ecology* 32:731–36
- Hobday AJ, Alexander LV, Perkins SE, Smale DA, Straub SC, et al. 2016. A hierarchical approach to defining marine heatwaves. *Prog. Oceanogr.* 141:227–38
- Hobday AJ, Oliver ECJ, Sen Gupta A, Benthuisen JA, Burrows MT, et al. 2018a. Categorizing and naming marine heatwaves. *Oceanography* 31(2):162–73
- Hobday AJ, Spillman CM, Eveson JP, Hartog JR, Zhang X, Brodie S. 2018b. A framework for combining seasonal forecasts and climate projections to aid risk management for fisheries and aquaculture. *Front. Mar. Sci.* 5:137
- Holbrook NJ, Claar DC, Hobday AJ, McInnes KL, Oliver ECJ, et al. 2020a. ENSO driven ocean extremes and ecosystem impacts. In *El Niño Southern Oscillation in a Changing Climate*, ed. MJ McPhaden, A Santoso, W Cai, pp. 409–28. New York: Wiley & Sons
- Holbrook NJ, Scannell HA, Gupta AS, Benthuisen JA, Feng M, et al. 2019. A global assessment of marine heatwaves and their drivers. *Nat. Commun.* 10:2624
- Holbrook NJ, Sen Gupta A, Oliver ECJ, Hobday AJ, Benthuisen JA, et al. 2020b. Keeping pace with marine heatwaves. *Nat. Rev. Earth Environ.* In press. <https://doi.org/10.1038/s43017-020-0068-4>
- Holt SA, Holt GJ. 1983. Cold death of fishes at Port Aransas, Texas: January 1982. *Southwest. Nat.* 28:464–66



- Hu D, Wu L, Cai W, Sen Gupta A, Ganachaud A, et al. 2015. Pacific western boundary currents and their roles in climate. *Nature* 522:299–308
- Hu Z, Kumar A, Jha B, Zhu J, Huang B. 2017. Persistence and predictions of the remarkable warm anomaly in the northeastern Pacific Ocean during 2014–16. *J. Clim.* 30:689–702
- Hughes TP, Kerry JT, Álvarez-Noriega M, Álvarez-Romero JG, Anderson KD, et al. 2017. Global warming and recurrent mass bleaching of corals. *Nature* 543:373–77
- Jacox MG. 2019. Marine heatwaves in a changing climate. *Nature* 571:485–87
- Jacox MG, Alexander MA, Siedlecki S, Chen K, Kwon Y-O, et al. 2020. Seasonal-to-interannual prediction of US coastal marine ecosystems: forecast methods, mechanisms of predictability, and priority developments. *Prog. Oceanogr.* 183:102307
- Jacox MG, Alexander MA, Stock CA, Hervieux G. 2017. On the skill of seasonal sea surface temperature forecasts in the California Current System and its connection to ENSO variability. *Clim. Dyn.* 53:7519–33
- Jacox MG, Tommasi D, Alexander MA, Hervieux G, Stock CA. 2019. Predicting the evolution of the 2014–2016 California Current System marine heatwave from an ensemble of coupled global climate forecasts. *Front. Mar. Sci.* 6:497
- Kataoka T, Tozuka T, Yamagata T. 2017. Generation and decay mechanisms of Ningaloo Niño/Niña. *J. Geophys. Res. Oceans* 122:8913–32
- Kay JE, Deser C, Phillips A, Mai A, Hannay C, et al. 2015. The Community Earth System Model (CESM) large ensemble project: a community resource for studying climate change in the presence of internal climate variability. *Bull. Am. Meteorol. Soc.* 96:1333–49
- Leadbetter MR, Lindgren G, Rootzén H. 1983. *Extremes and Related Properties of Random Sequences and Processes*. Berlin: Springer
- Lee T, Hobbs WR, Willis JK, Halkides D, Fukumori I, et al. 2010. Record warming in the South Pacific and western Antarctica associated with the strong central-Pacific El Niño in 2009–10. *Geophys. Res. Lett.* 37:L19704
- Lerorato JC, Nakamura Y. 2019. Unpredictable extreme cold events: a threat to range-shifting tropical reef fishes in temperate waters. *Mar. Biol.* 166:110
- Lewis SC, Mallela J. 2018. A multifactor risk analysis of the record 2016 Great Barrier Reef bleaching. *Bull. Am. Meteorol. Soc.* 99:S144–49
- Lim Y, Son S-W, Kim D. 2018. MJO prediction skill of the subseasonal-to-seasonal prediction models. *J. Clim.* 31:4075–94
- Lima FP, Wetthey DS. 2012. Three decades of high-resolution coastal sea surface temperatures reveal more than warming. *Nat. Commun.* 3:704
- Ling S, Johnson C, Ridgway K, Hobday A, Haddon M. 2009. Climate-driven range extension of a sea urchin: inferring future trends by analysis of recent population dynamics. *Glob. Change Biol.* 15:719–31
- Liu G, Heron SF, Eakin CM, Muller-Karger FE, Vega-Rodriguez M, et al. 2014. Reef-scale thermal stress monitoring of coral ecosystems: new 5-km global products from NOAA Coral Reef Watch. *Remote Sens.* 6:11579–606
- Manta G, de Mello S, Trinchin R, Badagian J, Barreiro M. 2018. The 2017 record marine heatwave in the southwestern Atlantic shelf. *Geophys. Res. Lett.* 45:12449–56
- Marshall AG, Hendon HH, Feng M, Schiller A. 2015. Initiation and amplification of the Ningaloo Niño. *Clim. Dyn.* 45:2367–85
- Mavrakīs AF, Tsiros IX. 2019. The abrupt increase in the Aegean sea surface temperature during June 2007 – a marine heatwave event? *Weather* 74:201–7
- McPhaden MJ, Zebiak SE, Glantz MH. 2006. ENSO as an integrating concept in earth science. *Science* 314:1740–45
- Mills KE, Pershing AJ, Brown CJ, Chen Y, Chiang F-S, et al. 2013. Fisheries management in a changing climate: lessons from the 2012 ocean heat wave in the Northwest Atlantic. *Oceanography* 26(2):191–95
- Moisan JR, Niiler PP. 1998. The seasonal heat budget of the North Pacific: net heat flux and heat storage rates (1950–1990). *J. Phys. Oceanogr.* 28:401–21
- Myers TA, Mechoso CR, Cesana GV, DeFlorio MJ, Waliser DE. 2018. Cloud feedback key to marine heatwave off Baja California. *Geophys. Res. Lett.* 45:4345–52



- Newman M, Wittenberg AT, Cheng L, Compo GP, Smith CA. 2018. The extreme 2015/16 El Niño, in the context of historical climate variability and change. *Bull. Am. Meteorol. Soc.* 99:S16–20
- Olita A, Sorgente R, Natale S, Gaberšek S, Ribotti A, et al. 2007. Effects of the 2003 European heatwave on the Central Mediterranean Sea: surface fluxes and the dynamical response. *Ocean Sci.* 3:273–89
- Oliver ECJ. 2019. Mean warming not variability drives marine heatwave trends. *Clim. Dyn.* 53:1653–59
- Oliver ECJ, Benthuisen JA, Bindoff NL, Hobday AJ, Holbrook NJ, et al. 2017. The unprecedented 2015/16 Tasman Sea marine heatwave. *Nat. Commun.* 8:16101
- Oliver ECJ, Burrows MT, Donat MG, Sen Gupta A, Alexander LV, et al. 2019. Projected marine heatwaves in the 21st century and the potential for ecological impact. *Front. Mar. Sci.* 6:734
- Oliver ECJ, Donat MG, Burrows MT, Moore PJ, Smale DA, et al. 2018a. Longer and more frequent marine heatwaves over the past century. *Nat. Commun.* 9:1324
- Oliver ECJ, Perkins-Kirkpatrick SE, Holbrook NJ, Bindoff NL. 2018b. Anthropogenic and natural influences on record 2016 marine heat waves. *Bull. Am. Meteorol. Soc.* 99:S44–48
- Oliver ECJ, Wotherspoon SJ, Chamberlain MA, Holbrook NJ. 2014a. Projected Tasman Sea extremes in sea surface temperature through the twenty-first century. *J. Clim.* 27:1980–98
- Oliver ECJ, Wotherspoon SJ, Holbrook NJ. 2014b. Estimating extremes from global ocean and climate models: a Bayesian hierarchical model approach. *Prog. Oceanogr.* 122:77–91
- Paz-Garca DA, Balart EF, Garca-de León FJ. 2012. Cold water bleaching of *Pocillopora* in the Gulf of California. In *Proceedings of 12th International Coral Reef Symposium*, ed. D Yellowlees, TP Hughes, pap. ICRS2012_9A_10. Townsville, Aust.: James Cook Univ.
- Pearce AF, Feng M. 2013. The rise and fall of the “marine heat wave” off Western Australia during the summer of 2010/2011. *J. Mar. Syst.* 111:139–56
- Pearce AF, Lenanton R, Jackson G, Moore J, Feng M, Gaughan D. 2011. *The “marine heat wave” off Western Australia during the summer of 2010/11*. Tech. Rep. 222, West. Aust. Fish. Mar. Res. Lab., North Beach, Aust.
- Perkins SE, Alexander LV. 2013. On the measurement of heat waves. *J. Clim.* 26:4500–17
- Perkins SE, Alexander LV, Nairn J. 2012. Increasing frequency, intensity and duration of observed global heatwaves and warm spells. *Geophys. Res. Lett.* 39:L20714
- Perkins-Kirkpatrick S, King A, Coughon E, Holbrook N, Grose M, et al. 2019. The role of natural variability and anthropogenic climate change in the 2017/18 Tasman Sea marine heatwave. *Bull. Am. Meteorol. Soc.* 100:S105–10
- Philander SGH. 1983. El Niño Southern Oscillation phenomena. *Nature* 302:295–301
- Pilo GS, Holbrook NJ, Kiss AE, Hogg AM. 2019. Sensitivity of marine heatwave metrics to ocean model resolution. *Geophys. Res. Lett.* 46:14604–12
- Power S, Delage F, Wang G, Smith I, Kociuba G. 2017. Apparent limitations in the ability of CMIP5 climate models to simulate recent multi-decadal change in surface temperature: implications for global temperature projections. *Clim. Dyn.* 49:53–69
- Rebert J-P, Donguy J-R, Eldin G, Wyrski K. 1985. Relations between sea level, thermocline depth, heat content, and dynamic height in the tropical Pacific Ocean. *J. Geophys. Res. Oceans* 90:11719–25
- Roberts H, Rouse L, Walker ND, Hudson J. 1982. Cold-water stress in Florida Bay and northern Bahamas: a product of winter cold-air outbreaks. *J. Sediment. Res.* 52:145–55
- Rouault M, Illig S, Bartholomae C, Reason C, Bentamy A. 2007. Propagation and origin of warm anomalies in the Angola Benguela upwelling system in 2001. *J. Mar. Syst.* 68:473–88
- Ruthrof KX, Breshears DD, Fontaine JB, Froend RH, Matusick G, et al. 2018. Subcontinental heat wave triggers terrestrial and marine, multi-taxa responses. *Sci. Rep.* 8:13094
- Salinger MJ, Renwick J, Behrens E, Mullan AB, Diamond HJ, et al. 2019. The unprecedented coupled ocean-atmosphere summer heatwave in the New Zealand region 2017/18: drivers, mechanisms and impacts. *Environ. Res. Lett.* 14:044023
- Schaeffer A, Roughan M. 2017. Subsurface intensification of marine heatwaves off southeastern Australia: the role of stratification and local winds. *Geophys. Res. Lett.* 44:5025–33
- Schlegel RW, Oliver ECJ, Hobday AJ, Smit AJ. 2019. Detecting marine heatwaves with sub-optimal data. *Front. Mar. Sci.* 6:737



- Schlegel RW, Oliver ECJ, Wernberg T, Smit AJ. 2017. Nearshore and offshore co-occurrence of marine heatwaves and cold-spells. *Prog. Oceanogr.* 151:189–205
- Sen Gupta A, McGregor S, Van Sebille E, Ganachaud A, Brown JN, Santoso A. 2016. Future changes to the Indonesian Throughflow and Pacific circulation: the differing role of wind and deep circulation changes. *Geophys. Res. Lett.* 43:1669–78
- Smale DA, Wernberg T, Oliver ECJ, Thomsen M, Harvey BP, et al. 2019. Marine heatwaves threaten global biodiversity and the provision of ecosystem services. *Nat. Clim. Change* 9:306–12
- Sparnocchia S, Schiano M, Picco P, Bozzano R, Cappelletti A. 2006. The anomalous warming of summer 2003 in the surface layer of the central Ligurian Sea (western Mediterranean). *Ann. Geophys.* 24:443–52
- Spillman C, Alves O, Hudson D. 2013. Predicting thermal stress for coral bleaching in the Great Barrier Reef using a coupled ocean–atmosphere seasonal forecast model. *Int. J. Climatol.* 33:1001–14
- Stott PA, Stone DA, Allen MR. 2004. Human contribution to the European heatwave of 2003. *Nature* 432:610–14
- Stuart-Smith RD, Edgar GJ, Barrett NS, Kininmonth SJ, Bates AE. 2015. Thermal biases and vulnerability to warming in the world's marine fauna. *Nature* 528:88–92
- Sugimoto S, Qiu B, Kojima A. 2020. Marked coastal warming off Tokai attributable to Kuroshio large meander. *J. Oceanogr.* 76:141–54
- Tan H, Cai R. 2018. What caused the record-breaking warming in East China Seas during August 2016? *Atmos. Sci. Lett.* 19:853
- Taschetto AS, Sen Gupta A, Jourdain NC, Santoso A, Ummenhofer CC, England MH. 2014. Cold tongue and warm pool ENSO events in CMIP5: mean state and future projections. *J. Clim.* 27:2861–85
- Tseng Y-H, Ding R, Huang X-M. 2017. The warm Blob in the northeast Pacific—the bridge leading to the 2015/16 El Niño. *Environ. Res. Lett.* 12:054019
- Tuckett CA, Wernberg T. 2018. High latitude corals tolerate severe cold spell. *Front. Mar. Sci.* 5:14
- Walsh JE, Thoman RL, Bhatt US, Bieniek PA, Brettschneider B, et al. 2018. The high latitude marine heat wave of 2016 and its impacts on Alaska. *Bull. Am. Meteorol. Soc.* 99:S39–43
- Weller E, Min S-K, Lee D, Kug J-S, Cai W, Yeh S-W. 2015. Human contribution to the 2014 record high sea surface temperatures over the western tropical and northeast Pacific Ocean. *Bull. Am. Meteorol. Soc.* 96:S100–4
- Wernberg T, Bennett S, Babcock RC, De Bettignies T, Cure K, et al. 2016. Climate-driven regime shift of a temperate marine ecosystem. *Science* 353:169–72
- Wernberg T, Smale DA, Tuya F, Thomsen MS, Langlois TJ, et al. 2013. An extreme climatic event alters marine ecosystem structure in a global biodiversity hotspot. *Nat. Clim. Change* 3:78–82
- WMO (World Meteorol. Organ.). 2017. *WMO guidelines on the calculation of climate normals*. Doc. WMO-No. 1203, WMO, Geneva
- WMO (World Meteorol. Organ.). 2018. *Guide to climatological practices*. Doc. WMO-No. 100, WMO, Geneva
- Wu L, Cai W, Zhang L, Nakamura H, Timmermann A, et al. 2012. Enhanced warming over the global subtropical western boundary currents. *Nat. Clim. Change* 2:161–66
- Zhang N, Feng M, Hendon HH, Hobday AJ, Zinke J. 2017. Opposite polarities of ENSO drive distinct patterns of coral bleaching potentials in the southeast Indian Ocean. *Sci. Rep.* 7:2443
- Zhao S, Jin F-F, Stuecker MF. 2019. Improved predictability of the Indian Ocean Dipole using seasonally modulated ENSO forcing forecasts. *Geophys. Res. Lett.* 46:9980–90

RELATED RESOURCES

- Australian Bureau of Meteorology ocean temperature outlooks: <http://www.bom.gov.au/oceanography/oceantemp/sst-outlook-map.shtml>
- California Current Marine Heatwave Tracker: <https://www.integratedecosystemassessment.noaa.gov/regions/california-current/cc-projects-blobtracker>
- Climate Prediction Center seasonal climate forecast from CFSv2: <https://www.cpc.ncep.noaa.gov/products/CFSv2/CFSv2seasonal.shtml>



Coral Reef Watch satellite monitoring and modeled outlooks: <https://coralreefwatch.noaa.gov/satellite>
Fishforecasts Marine Heatwave Monitoring: <https://fishforecasts.dtu.dk/heatwaves>
Integrated Marine Observing System OceanCurrent: <http://oceancurrent.imos.org.au>
Marine Heatwaves International Working Group Marine Heatwave Tracker: <http://www.marineheatwaves.org/tracker.html>
T-MEDNet Mediterranean Marine Heatwaves: <http://t-mednet.org/t-resources/marine-heatwaves>

